The background of the slide is a photograph of a rural landscape. In the foreground, there is a lush green field, possibly a vineyard or a crop field. Behind it, there are rolling hills with dry, golden-brown grass. In the distance, more hills are visible under a clear blue sky. The text is overlaid on this image.

Reporting Biomonitoring Results to Individuals and Medical and Public Audiences: Challenges and Opportunities

Asa Bradman, PhD

September 25, 2007


*UC Berkeley Center for Children's Environmental
Health Research*

Outline

- What is CHAMACOS?
- Why do it
- What have we found?
- Case studies



Berkeley
University of California



A Community-University Partnership




South County Outreach Effort



Clinica de Salud del Valle de Salinas







Results of Community Assessment

The Community has an interest in Environmental Health Research.

“There is a need to know how the environment is affecting us.”

Children’s Health is a priority concern in the Community.

“I think you will get a very positive response from the women. They are very interested in their children’s health and how to improve it. You need to give them access to their results.”

The Center must share results with the Community.

“I think the reaction of the community is going to be positive but you guys need to keep them informed about the study and talk the way you’ve talked to me about the community’s long term benefits from the study, and plans to stay in the community for further studies.”

Research must be culturally sensitive.

“Hire people who can work effectively with our community, people who understand the culture.”



Objectives


- Assess exposures to pregnant women and children.
- Determine relationship with :
 - neurodevelopment
 - growth
 - respiratory disease
- Reduce exposures to children and pregnant women with interventions and community outreach.
- Inform policies to reduce the incidence of environmentally-related disease



	1 st Tri	2 nd Tri	Deliver y	6 M	1 Y	2 Y	3½ Y	5 Y	7 Y
• Maternal Urine	✓	✓	✓	✓					
• Paternal Urine			✓						
• Maternal Blood		✓	✓						✓
• Cord Blood			✓						
• Breast Milk			✓	✓					
• Child Urine				✓	✓	✓	✓	✓	
• Child Blood					✓	✓		✓	✓
• Child Saliva							✓	✓	




Prenatal and Child Total Dialkyl Phosphate (DAP) Metabolites





Maternal Exposure to OCs

Comparison of serum organochlorine (OC) levels in the CHAMACOS Cohort and NHANES 1999-2000 populations
(All values are nanogram/gram lipid)




Bradman et al 2007.



OPs and Neonatal Neurodevelopment

Percentage of Neonates with >3 Abnormal Reflexes
(Brazelton Neonatal Behavior Assessment Scale)



$$X^2_{\text{trend}} = 6.7$$



Neurodevelopment at 2 years

OPs

	<u>Bayley MDI</u>	<u>CBCL PDD</u>
	<u>β (95% CI)</u>	<u>OR (95% CI)</u>
Prenatal OPs	-3.5 (-6.6, -0.5)	2.3 (1.0, 5.2)

Home > ForbesLife > Health

DDT Linked to Developmental Delays in Babies

07.05

WED
sugg
were

DDT
consi

The n
malai
Brend
at the

Envir
and it
it's be

While
on hu
in the
branc

E-mail | Print | Comments | E-Mail Newsletters | My Yahoo! | RSS

E-Mail Alerts
Get new stories by e-mail as they are published FREE




LEARN ABOUT TREATMENT >>>

Learn About Treatment >>>

Site Search

Home > Kids > Child Development



Visit Our Health Centers
* All Health Centers *
Acne
Addictions
AIDS/HIV
Alcohol Abuse

News
Health Encyclopedia
In-Depth Reports
Check a Symptom
Procedures
Animations Library
Video Library
Dr. Koop's Health Tools
Drug Information
Natural Medicine

What's HOT
Joint Pain Explained
GERD
Treatment Options for Enlarged Prostate

Subscribe to our Newsletter
your email address Go

contact us
privacy policy

Printer Friendly
DDT Linked to Developmental Delays in Babies
Breast-feeding seems to offset some of the pesticide's negative effects, study finds

By Randy Dottinga
HealthDay Reporter



exposed to the pesticide.

WEDNESDAY, July 5 (HealthDay News) -- New research out of California suggests that DDT causes developmental delays in infants whose mothers were

DDT is currently banned in the United States, but officials in Africa are considering expanding its use to combat mosquitoes that spread malaria.

The new findings -- along with the potential benefits of DDT use to reduce malaria -- "need to be considered by the study author Brenda Eskenazi, a professor of maternal and child health at the University of California, Berkeley."

[Text continued below](#)

Los Angeles Times
latimes.com

search Go

4:48 PM PDT, July 6, 2006

Sponsored by
CEDARS-SINAI

- HOME
- Site Map
- Health
- Medicine
- Fitness
- Nutrition
- Columns
- NEWS
- California | Local
- National
- World
- Entertainment News
- Business
- Sports
- Politics
- Opinion
- Columnists
- Print Edition
- RSS What is RSS?
- Calendar/ive
- Travel
- West Magazine
- House & Garden
- Health
- Food
- Car Culture
- Obituaries
- Crossword, Sudoku
- All Sections
- Corrections
- BUY, SELL & MORE
- Jobs

Health : Medicine

DDT Study Finds New Hazard to Young Children

Babies born in the U.S. to mothers emigrating from Mexico show mental and physical impairment, a UC Berkeley survey finds.

By Marla Cone, Times Staff Writer
July 5, 2006

Babies and toddlers of California farmworkers exposed to the insecticide DDT have neurological effects that are severe enough in some cases to slow their mental and physical development, according to research by UC Berkeley scientists published today.

The federally funded research involving the children of women who recently emigrated from Mexico to the Salinas Valley is the first in the United States to indicate that the pesticide harms human brain development.

ADVERTISEMENT

Nurses:
Apply your skills in a world-class medical environment

Search here

CEDARS-SINAI
LEADING THE QUEST FOR HEALTH.

"This suggests that ... DDT has effects that no one even thought to test for back when it was in use," said Dr. Walter Rogan, an epidemiologist with the National Institute of Environmental Health Sciences. He was not involved in the new study, published in the journal Pediatrics.

Because DDT was banned more than 30 years ago in the United States and most developed countries, the findings have particular relevance for the ongoing, controversial use in Africa to combat malaria.

UC Berkeley scientists measured levels of various pesticides in 360 pregnant women, nearly all of whom were born in Mexico, and tested the mental and motor skills of their infants and toddlers, who were born in the Salinas Valley.

bbc.co.uk

Home TV Radio Talk Where I Live A-Z Index



BBC NEWS

UK version International version About the versions | Low graphics | Help | Co

News Front Page

- World
- UK
- England
- Northern Ireland
- Scotland
- Wales
- Business
- Politics
- Health
- Medical notes
- Education
- Science/Nature
- Technology

Last Updated: Wednesday, 5 July 2006, 04:08 GMT 05:08 UK

E-mail this to a friend

Printable version

DDT 'link' to slow child progress

Children exposed to the pesticide DDT while in the womb experience development problems, researchers say.



The pesticide was banned in the US and UK in the 1970s, but it is still used in some countries to kill malaria-carrying mosquitoes.

It was already known DDT was

DDT is used to protect against

SEE ALSO
DDT pr
27 Jun
DDT lin
12 Jul
Premat
16 May

RELATED
Pediatr
Univers
MRC Hu
Science
The BBC
content

TOP HEA
Painkill
Therap
cancer
Charts

RSS

MOST PO
MOST E

- 1 Comic
 - 2 Video
 - 3 Putin
 - 4 Gaza
 - 5 Go-ah
- extrac

MO
- HI
- AI
- In
- DE
- Ev
- M
- M
MO
- HI
- AI
- Ed
- Br
RE
- Bi
- Tr
- Fa
- M
- Hi
- Ki
- Po

CO



say their
in mind when

ent long after
issues of animals


have similar

US, DDT is still
ODE is

els of DDT and
ant women, the




Distribution of Average Cumulative Pesticide Dose Estimates for CHAMACOS Women





PON1 Activity Varies by Age and Genotype



- PON1 newborns lower than adults;


therefore, greater susceptibility to OP's

- Children with QQ genotype have low level of enzyme;

therefore, may be at highest risk



Median DM Metabolite Levels by Fruit and Vegetable consumption





CHAMACOS Biomonitoring Uses

- Characterize exposure
- Evaluate exposure trends
- Evaluate health effects
- Estimate risks



Case Studies in Communication

- **IRB education**
- **Community involvement**
- **Challenges by the medical community.**
- **Final approach.**
- **Technical challenges**
- **General community**



UC Berkeley CPHS

Forbidden to return individual results (1999)



Community Engagement

- CPHS agreed to reevaluate decision.
- Convened meetings with wide range of community and research partners.



Community Engagement

- Doctors strongly objected to returning individual results –burden of interpreting non-clinical tests.
- Advocates strongly supported returning individual results – right to know.
- Industry supported returning individual results – individual right.



Final approach accepted by IRB

- Participants can request results (opt in).
- Participants informed:
 - In person;
 - Results placed in context of population and reference data (NHANES);
 - Emphasize research context;
 - Offer follow-up testing;
 - Don't stigmatize;
 - If known risk (e.g., lead), follow guidelines.
 - Only ~20 have asked for results;
- Secondary protocol to re-test “high” results.



History Repeats Itself: CPHS II

CPHS II: 12/8/06

“test results for individual subjects ... not ...meaningful and ... likely to cause unnecessary alarm ... members felt strongly that... subjects ... only be provided with a copy of the manuscript.”



Technical Challenges

Biomarkers may not reflect exposure.

For example, DAPs in the environment and food suggest urinary OP biomarkers require additional interpretation.



Technical Challenges

For many non-persistent compounds, good intra-individual correlation over a day. Poor over several days.



Correlation of DAP metabolites in spot urine samples collected 1-6 days apart

Days between collection
(n)

1	2	3	4	5	6
(288)	(241)	(210)	(144)	(72)	(25)

Total DAPS	0.34**	0.21**	0.12	0.03	0.13	-0.05
Dimethyls	0.38**	0.27**	0.17*	0.07	0.16	-0.09
Diethyls	0.18**	0.03	0.00	0.02	0.05	-0.22

**p<0.01

*p<0.05



Correlation of 24 hr samples collected three days apart (n=25)

Total DAPS 0.11


Dimethyls 0.11

Diethyls 0.06

$p > 0.6$



24 hr urine samples: Within and between individual variability



Estimated within and between variability (SD)

Between 0.36

Within 1.01



Technical Challenges

- Individual sampling results for some non-persistent chemicals may be meaningless.
- Whereas population exposure range is well characterized.



Reporting results to the General Community



Research partners and advisory boards:

- All findings presented during in-person meetings;
- Submitted journal articles provided for comment;
- Press releases distributed for comment;
- Review is advisory only;



General community:

- Frequent participant fora.
- Presentations to broad cross-section of community:
 - Migrant education;
 - Community groups
 - Churches;
 - Local governments;
 - Grower organizations



Community Outreach





Outreach and translation

In the home.



In the community.



In the fields.





Investigators

Exposure Studies

- Asa Bradman
- Tom McKone
- Dana Barr CDC
- Rosemary Castorina
- Martha Harnly, DHS
- Jim Leckie, Stanford
- Marcia Nishioka, Batelle
- Lesliam Quiros

Health Studies

- Brenda Eskenazi
- Ira Tager
- Kim Harley
- Laura Fenster, DHS
- Caroline Johnson
- Michael Lipsett, DHS
- Janet Macher, DHS

Mechanism Studies Center

- Nina Holland
- John Casida

Biostatistical Core

- Nick Jewell
- Alan Hubbard
- Amy Marks

Intervention and Community Outreach

- Abbey Alkon, UCSF
- Lisa Goldman
- Alicia Salvatore
- Celina Trujillo

Clinica de Salud

- John Silva
- Max Cuevas

Natividad Medical

• Marc Tunzi





- **California Wellness Foundation**
- **Switzer Foundation**
- **UC Mexus**